



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\cot x = \frac{1}{x} - \frac{1}{\pi - x} + \frac{1}{\pi + x} - \frac{1}{2\pi - x} + \frac{1}{2\pi + x} - \dots^*$$

$$= \frac{1}{x} + \frac{x^2}{\pi - 2x} + \frac{(\pi - x)^2}{2x} + \frac{(\pi + x)^2}{(\pi - 2x)} + \dots$$

$$\sec x = \frac{4\pi}{\pi^2 - 4x^2} - \frac{3.4\pi}{3^2\pi^2 - 4x^2} + \frac{4.5\pi}{5^2\pi^2 - 4x^2} - \frac{4.7\pi}{7^2\pi^2 - 4x^2} + \frac{4.9\pi}{9^2\pi^2 - 4x^2} - \dots^\dagger$$

$$= \frac{4\pi}{\pi^2 - 4x^2} + \frac{3(\pi^2 - 4x^2)^2}{6\pi^2 + 8x^2} + \frac{(9\pi^2 - 4x^2)^2}{30\pi^2 + 8x^2} + \dots$$

$$\csc x = \frac{1}{x} + \frac{2x}{\pi^2 - x^2} - \frac{2x}{2^2\pi^2 - x^2} + \frac{2x}{3^2\pi^2 - x^2} - \frac{2x}{4^2\pi^2 - x^2} + \dots$$

$$= a - \frac{ax - 1}{x} + \frac{2x}{\pi^2 - x^2} - \frac{2x}{4\pi^2 - x^2} + \dots^\ddagger$$

$$= \frac{a}{1} + \frac{ax - 1}{1} + \frac{2ax^3}{a\pi^2x - \pi^2 - x^2 - ax^3} + \frac{2x(ax - 2)(\pi^2 - x^2)^2}{6\pi^2} + \dots$$

Also solved by G. W. Greenwood.

268. Proposed by O. E. GLENN, Ph. D., Philadelphia, Pa.

Express the hyperbolic functions of x in the form of infinite continued fractions.

Solution by J. SCHEFFER, A. M., Professor of Mathematics, Kee Mar College, Hagerstown, Md.

For the conversion of $\tanh x$ and $\tanh x$, Legendre has given a very elegant solution.

Putting $f(x) = 1 + \frac{a}{x} + \frac{1}{2} \cdot \frac{a^2}{x(x+1)} + \frac{1}{2 \cdot 3} \cdot \frac{a^3}{x(x+1)(x+2)} + \dots$ we have

$$f(x+1) = 1 + \frac{a}{x+1} + \frac{1}{2} \cdot \frac{a^2}{(x+1)(x+2)} + \frac{1}{2 \cdot 3} \cdot \frac{a^3}{(x+1)(x+2)(x+3)} + \dots$$

$$\therefore f(x) - f(x+1) = \frac{a}{x(x+1)} + \frac{a^2}{x(x+1)(x+2)} + \frac{1}{2} \frac{a^3}{x(x+1)(x+2)(x+3)} + \dots$$

**Encyclopedia Britannica*, Vol. XXIII, p. 572.

†Euler's *Institu. Calc. Diff.*, II, 8, No. 223—224.

‡The quantity a may have any integral value which will avoid negative terms in the continued fraction.

$$= \frac{a}{x(x+1)} \left[1 + \frac{a}{x+2} + \frac{1}{2} \frac{a^2}{(x+2)(x+3)} + \dots \right] = \frac{a}{x(x+1)} f(x+2);$$

whence, putting $F(x) = \frac{a}{x} \cdot \frac{f(x+1)}{f(x)}$, $F(x+1) = \frac{a}{x+1} \frac{f(x+2)}{f(x+1)}$, we obtain

$$\frac{x}{a} \frac{f(x)}{f(x+1)} - \frac{x}{a} = \frac{1}{x+1} \cdot \frac{f(x+2)}{f(x+1)}, \quad \frac{1}{F(x)} - \frac{x}{a} = \frac{F(x+1)}{a}, \quad F(x) = \frac{a}{x + F(x+1)}$$

$$\therefore F(x+1) = \frac{a}{x+1+F(x+1)}, \quad F(x+2) = \frac{a}{x+2+F(x+2)}, \text{ etc.}$$

$$\therefore F(x) = \frac{a}{x} + \frac{a}{x+1} + \frac{a}{x+2} + \frac{a}{x+3} + \dots,$$

$$\text{where } F(x) = \frac{a}{x} \frac{1 + \frac{a}{x+1} + \frac{1}{2} \frac{a^2}{(x+1)(x+2)} + \frac{1}{2 \cdot 3} \frac{a^3}{(x+1)(x+2)(x+3)} + \dots}{1 + \frac{a}{x} + \frac{1}{2} \frac{a^2}{x(x+1)} + \frac{1}{2 \cdot 3} \frac{a^3}{x(x+1)(x+2)} + \dots}$$

$$\text{For } x = \frac{1}{2}, F(x) = 2a \cdot \frac{1 + \frac{4a}{2 \cdot 3} + \frac{16a^2}{2 \cdot 3 \cdot 4 \cdot 5} + \frac{64a^3}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6} + \dots}{1 + \frac{4a}{2} + \frac{16a^2}{2 \cdot 3 \cdot 4} + \frac{64a^3}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6} + \dots} = \frac{\epsilon^{2\sqrt{a}} - \epsilon^{-2\sqrt{a}}}{\epsilon^{2\sqrt{a}} + \epsilon^{-2\sqrt{a}}} \sqrt{a}.$$

$$\therefore \frac{\epsilon^{2\sqrt{a}} - \epsilon^{-2\sqrt{a}}}{\epsilon^{2\sqrt{a}} + \epsilon^{-2\sqrt{a}}} 2\sqrt{a} = \frac{4a}{1} + \frac{4a}{4} + \frac{4a}{5} + \dots$$

$$\text{Putting } 2\sqrt{a} = x, \quad \frac{e^x - e^{-x}}{e^x + e^{-x}} = \tanh x = \frac{x}{1} + \frac{x^2}{3} + \frac{x^2}{5} + \frac{x^3}{7} + \dots$$

$$\text{Also } \frac{e^{ix} - e^{-ix}}{e^{ix} + e^{-ix}} = \frac{ix}{1} - \frac{x^2}{3} - \frac{x^2}{5} - \dots$$

$$\therefore i \tan x = \frac{ix}{1} - \frac{x^2}{3} - \frac{x^2}{5} - \frac{ix^3}{7} - \dots \quad \text{or, } \tan x = \frac{x}{1} - \frac{x^2}{3} - \frac{x^2}{5} - \dots$$

a very simple and remarkable continued fraction.

Since $\sinh x = -i \sin ix$, and $\cosh x = \cos ix$, we get, by substituting ix for x in the continued fractions for $\sin x$ and $\cos x$,

$$\sinh x = \frac{x}{1} - \frac{x^3}{2.3+x^2} - \frac{2.3x^2}{4.5+x^2} - \frac{4.5x^2}{6.7+x^2} - \text{etc.},$$

$$\cosh x = \frac{1}{1} - \frac{1}{2+x^2} - \frac{2x^2}{3.4+x^2} - \frac{3.4x^2}{5.6+x^2} - \text{etc.}$$

GEOMETRY.

289 (Incorrectly numbered 288). Proposed by C. N. SCHMALL, College of the City of New York.

From a point P on a given circle to draw two chords such that, (α) chord $PA : \text{chord } PB = m : n$ (a given ratio), and, (β) arc $PA : \text{arc } PB = 1 : 3$.

Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Let O be the center of the circle, radius r . Also let angle $POA = 2\theta$, angle $POB = 6\theta$. Then $PA = 2r \sin \theta$, $PB = 2r \sin 3\theta$.

$$\frac{PA}{PB} = \frac{\sin \theta}{\sin 3\theta} = \frac{1}{3 - 4 \sin^2 \theta} = \frac{m}{n}. \quad \therefore \sin \theta = \frac{1}{2} \sqrt{\frac{3m-n}{m}}, \quad \sin 3\theta = \frac{n}{2m} \sqrt{\frac{3m-n}{m}}.$$

$$\therefore \text{Make angle } POA = 2 \sin^{-1} \frac{1}{2} \sqrt{\frac{3m-n}{m}}; \text{ angle } POB = 2 \sin^{-1} \frac{n}{2m} \sqrt{\frac{3m-n}{m}}.$$

Then chord $PA : \text{chord } PB = m : n$; arc $PA : \text{arc } PB = 1 : 3$.

290 (Incorrectly numbered 289). Proposed by J. J. QUINN, Ph. D., Scottdale, Pa.

(a) Suppose a circle described around the origin. Then at the end of a uniformly revolving radius r , a line equal to the diameter is pivoted. Find the equation of the locus of its extremity, if for every unit of angle its projection on the X axis is a constant linear unit, being the same part of the diameter as the angle is of π radians.

(b) Show how it can be applied to the trisection or multisection of an angle.

No solution has been received.

292 (Incorrectly numbered 290). Proposed by DR. L. E. DICKSON, The University of Chicago, Chicago, Ill.

Given nine points lying by threes in three columns and in three rows, draw through them, by continuous motion, a broken line composed of only four straight segments, and passing but once through each of the nine points. [A current puzzle.]